


Doc
W2
A2
9212P

(DOCUMENT SECTION)

PROJECT REPORT COMMITTEE ON FOOD RESEARCH U.S. QUARTERMASTER FOOD AND CONTAINER INSTITUTE FOR THE ARMED FORCES CHICAGO ILLINOIS		RESEARCH AND DEVELOPMENT BRANCH MILITARY PLANNING DIVISION OFFICE OF THE QUARTERMASTER GENERAL	
COOPERATING INSTITUTION University of Texas		LOCALITY Austin, Texas	
DIVISION Arts and Sciences		DEPARTMENT Bacteriology	
OFFICIAL INVESTIGATOR J. W. Foster		COLLABORATORS E. S. Wynne	
REPORT NO. 6		FILE NO. M-605	CONTRACT NO. W11-009-qm-70190
FOR PERIOD COVERING 1 July 1947 - 31 December 1947		INITIATION DATE 1 July 1946	
TITLE: [] PROGRESS REPORT [X] PHASE REPORT [] ANNUAL REPORT [] TERMINATION REPORT Spore formation and spore germination of anaerobic food spoilage organisms, especially <i>Clostridium botulinum</i> .			
SUMMARY			
<p>Data obtained in this study, together with information in the literature lead to the conclusion that the classical "dormancy" observed in anaerobic as well as aerobic spores is not a biological character of the spore itself, but is rather a function of the medium. "Dormancy" may be considered as a manifestation of the presence of anti-germination substances in normal media. "Dormancy" can be eliminated with varying degrees of success, depending on the medium and the nature of the inhibitive substances by the incorporation of various supplements, in particular soluble starch (0.1 percent), which absorbs some, but not all, the inhibitors.</p> <p>The term "dormancy" would appear to be of dubious value in connection with bacterial spores.</p>			
			
M-605 #6		- 1 -	
		Continued	

The Problem of "Dormancy" in Bacterial Spores¹

Jackson W. Foster and E. Staten Wynne²

Department of Bacteriology, University of Texas

Our experience with germination of spores of various species of Clostridia, especially C. botulinum, has led to conclusions which we believe warrant the consideration of investigators of this group of organisms and of the aerobic spore formers as well. Pertinent observations have been made by previous authors, but it appears that all the recent available evidence and various aspects of the dormancy problem have not hitherto been crystallized into one comprehensive concept.

The essential points already established (Morrison and Rettger, 1930a and 1930b for review of previous literature; Curran and Evans, 1937; Olsen and Scott, 1946; Wynne and Foster, 1948a; Foster and Wynne, 1948) are these:

(1) Colony counts from a spore suspension may be submaximum in a few days incubation, and increase with time over prolonged periods. This indicates delayed germination and is a well-known phenomenon for both aerobic and anaerobic spores. This is the characteristic feature of the classical dormancy.

(2) This effect depends on the medium used, it being marked in some, and non-existent in others.

1 The research which this paper reports was undertaken in cooperation with the Committee on Food Research of the Quartermaster Food and Container Institute for the Armed Forces. The views or conclusions contained in this report are those of the authors. They are not to be construed as necessarily reflecting the views or endorsement of the War Department.

2 Present address: Dept. of Plant Sciences, University of Oklahoma, Norman, Oklahoma.

(3) The final spore counts of a given suspension depend greatly upon the medium employed, though in each medium counts are at the maximum obtainable in it.

(4) Certain substances, notably soluble starch (0.1 per cent), eliminate the delayed germination in some media. In others the starch increases the final maximum count, whereas in still others it has no apparent effect. In general, all media containing complex nitrogenous materials preferred for counting spores, show the starch effect.

Data exemplifying all these points may be found in Table 1. From the information in the literature, as well as in Table 1, the following points may be made:

(a) The so-called "dormancy" in these organisms is not an inherent function of the spores, but rather of the medium in which the spores are placed to germinate.

(b) Even where "dormancy" occurred, or in media with low final counts, the large size of colonies speaks against a nutritional inadequacy.

(c) Accordingly, a plausible conclusion is that "dormancy" i.e., delayed germination, and also submaximal counts, are due to substances in normal media inhibitory to spore germination. "Dormancy" would then be only a manifestation of anti-germination substances in the medium and seemingly is, therefore, merely an artifact.

(d) In the case of the characteristic delayed rate of spore germination during "dormancy" one may consider this to be a function of sub-inhibitory concentrations of certain substances in the normal medium.

(e) In the case of submaximal counts, the inhibitory substances are of such nature and concentration that the germination rate of a portion of the spore population is zero, i.e., they never germinate, for all practical purposes.

(f) The inhibitors evidently are of different types, but one is the C_{18} unsaturated fatty acids (oleic, linoleic, linolenic) which are present in almost

(g) The appreciable increase in germination rate, occasioned by the soluble starch in some media, is due to absorption of certain inhibitory substances, including the C_{18} fatty acids. In some media starch has little or no effect in enhancing counts known to be low.

(h) The different degrees of inactivation of germination inhibitors by starch indicate that different media contain varying amounts of different inhibitors. Solvent extraction of some media indicates some of these naturally occurring inhibitors are non-lipid.

(i) The significance of these substances inhibitory to germination is evidenced by the fact that the fresh infusions of beef, pork, and liver, ordinarily considered ideal nutrients, contain significant amounts of germination inhibitors, as demonstrated by the considerable enhancement of germination caused by addition of starch. Incidentally, fresh beef or liver infusion with starch is just as suitable as the widely used (for anaerobes) pork infusion medium of Yesair, with added starch.

(j) Effects similar to that of starch have been observed in the past when supplements containing native proteins and polysaccharides are added to the usual bacteriological media. Though this action has been arbitrarily ascribed to unspecified nutritional factors, an equally plausible idea is that they function physically in the manner of starch, through colloid absorption of inhibitory substances, and in some cases this has been proved definitely.

(k) Unless shown that no effect is obtained by starch, the above data and ideas make it imperative that this substance be an invariable component of all media in work on anaerobic as well as aerobic spores where maximum counts in the shortest time are desired. Even though starch apparently does not counteract all known cases of retarded germination, it is the simplest overall treatment available without altering the nutritional status of the basal medium.

(1) The term "dormancy" in the above connection no longer has the degree of validity it formerly enjoyed, and there now appears to be some question as to the justification of its continued use, unless in cases where the effect is shown to be independent of the germination medium.

	Days	1	2	3	4
Yeast extract 0.2 percent	-	45	42	38	-
Yeast extract 0.1 percent	-	45	42	38	-
Yeast extract 0.05 percent	-	45	42	38	-
Yeast extract 0.025 percent	-	45	42	38	-
Yeast extract 0.0125 percent	-	45	42	38	-
Yeast extract 0.00625 percent	-	45	42	38	-
Yeast extract 0.003125 percent	-	45	42	38	-
Yeast extract 0.0015625 percent	-	45	42	38	-
Yeast extract 0.00078125 percent	-	45	42	38	-
Yeast extract 0.000390625 percent	-	45	42	38	-
Yeast extract 0.0001953125 percent	-	45	42	38	-
Yeast extract 0.00009765625 percent	-	45	42	38	-
Yeast extract 0.000048828125 percent	-	45	42	38	-
Yeast extract 0.0000244140625 percent	-	45	42	38	-
Yeast extract 0.00001220703125 percent	-	45	42	38	-
Yeast extract 0.000006103515625 percent	-	45	42	38	-
Yeast extract 0.0000030517578125 percent	-	45	42	38	-
Yeast extract 0.00000152587890625 percent	-	45	42	38	-
Yeast extract 0.000000762939453125 percent	-	45	42	38	-
Yeast extract 0.0000003814697265625 percent	-	45	42	38	-
Yeast extract 0.00000019073486328125 percent	-	45	42	38	-
Yeast extract 0.000000095367431640625 percent	-	45	42	38	-
Yeast extract 0.0000000476837158203125 percent	-	45	42	38	-
Yeast extract 0.00000002384185791015625 percent	-	45	42	38	-
Yeast extract 0.000000011920928955078125 percent	-	45	42	38	-
Yeast extract 0.0000000059604644775390625 percent	-	45	42	38	-
Yeast extract 0.00000000298023223876953125 percent	-	45	42	38	-
Yeast extract 0.000000001490116119384765625 percent	-	45	42	38	-
Yeast extract 0.0000000007450580596923828125 percent	-	45	42	38	-
Yeast extract 0.00000000037252902984619140625 percent	-	45	42	38	-
Yeast extract 0.000000000186264514923095703125 percent	-	45	42	38	-
Yeast extract 0.0000000000931322574615478515625 percent	-	45	42	38	-
Yeast extract 0.00000000004656612873077392578125 percent	-	45	42	38	-
Yeast extract 0.000000000023283064365386962890625 percent	-	45	42	38	-
Yeast extract 0.0000000000116415321826934814453125 percent	-	45	42	38	-
Yeast extract 0.00000000000582076609134674072265625 percent	-	45	42	38	-
Yeast extract 0.000000000002910383045673370361328125 percent	-	45	42	38	-
Yeast extract 0.0000000000014551915228366851806640625 percent	-	45	42	38	-
Yeast extract 0.00000000000072759576141834259033203125 percent	-	45	42	38	-
Yeast extract 0.000000000000363797880709171295166015625 percent	-	45	42	38	-
Yeast extract 0.0000000000001818989403545856475830078125 percent	-	45	42	38	-
Yeast extract 0.00000000000009094947017729282379150390625 percent	-	45	42	38	-
Yeast extract 0.000000000000045474735088646411895751953125 percent	-	45	42	38	-
Yeast extract 0.0000000000000227373675443232059478759765625 percent	-	45	42	38	-
Yeast extract 0.00000000000001136868377216160297393798828125 percent	-	45	42	38	-
Yeast extract 0.000000000000005684341886080801486968994140625 percent	-	45	42	38	-
Yeast extract 0.0000000000000028421709430404007434844970703125 percent	-	45	42	38	-
Yeast extract 0.00000000000000142108547152020037174224853515625 percent	-	45	42	38	-
Yeast extract 0.000000000000000710542735760100185871124267578125 percent	-	45	42	38	-
Yeast extract 0.0000000000000003552713678800500929355621337890625 percent	-	45	42	38	-
Yeast extract 0.00000000000000017763568394002504646778106689453125 percent	-	45	42	38	-
Yeast extract 0.000000000000000088817841970012523233890533447265625 percent	-	45	42	38	-
Yeast extract 0.0000000000000000444089209850062616169452667236328125 percent	-	45	42	38	-
Yeast extract 0.00000000000000002220446049250313080847263336181640625 percent	-	45	42	38	-
Yeast extract 0.000000000000000011102230246251565404236316680908203125 percent	-	45	42	38	-
Yeast extract 0.0000000000000000055511151231257827021118158404541015625 percent	-	45	42	38	-
Yeast extract 0.00000000000000000277555756156289135105590792022705078125 percent	-	45	42	38	-
Yeast extract 0.000000000000000001387778780781445675527953960113525390625 percent	-	45	42	38	-
Yeast extract 0.0000000000000000006938893903907228377639769800567626953125 percent	-	45	42	38	-
Yeast extract 0.00000000000000000034694469519536141888198849002838134765625 percent	-	45	42	38	-
Yeast extract 0.000000000000000000173472347597680709440994245014190673828125 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000867361737988403547204971225070953369140625 percent	-	45	42	38	-
Yeast extract 0.00000000000000000004336808689942017736024856125354766845703125 percent	-	45	42	38	-
Yeast extract 0.000000000000000000021684043449710088680124280626773834228515625 percent	-	45	42	38	-
Yeast extract 0.000000000000000000010842021724855044340062140313386917114278125 percent	-	45	42	38	-
Yeast extract 0.000000000000000000005421010862427522170031070156694585571390625 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000027105054312137610850155350783472927856953125 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000135525271560688054250776753917364639284765625 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000677626357803440271253883769586823196423828125 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000003388131789017201356269418847934115982119140625 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000016940658945086006781347094239670579910595703125 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000084703294725430033906735471198352899552978515625 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000423516473627150169533677355991764497764892578125 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000002117582368135750847668386779958822488824462890625 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000010587911840678754238341933899794112444122314453125 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000052939559203393771191709669498970562220611572265625 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000264697796016968855958548347494853111030557861328125 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000001323488980084844279792741737474265555152789306640625 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000006617444900424221398963708687371327775763946533203125 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000033087224502121106994818543436856638878819732666015625 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000000165436122510605534974092717184283194394098663330078125 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000000827180612553027674870463585921415971970493316650390625 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000004135903062765138374352317929607079859852465583251953125 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000000020679515313825691871761589648035399299262327916259765625 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000000103397576569128459358807948240176996496311639581298828125 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000000516987882845642296794039741200884982481558197906494140625 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000000002584939414228211483970198706004424912407790989532470703125 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000000012924697071141057419850993530022124562038954947654853515625 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000000064623485355705287099254967650110622810194774737792767578125 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000000000323117426778526435496274838250553114050973873688963837890625 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000000001615587133892632177481374191252765570254869368444819189453125 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000000008077935669463160887406870956263827851274346842224095947265625 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000000000040389678347315804437034354781319139256371734211120479736328125 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000000000201948391736579022185171773906595696281858671055602398681640625 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000000001009741958682895110925858869532978481409293355278011993408203125 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000000000005048709793414475554629294347664892407046466776390059967041015625 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000000000025243548967072377773146471738324462035232333881950299835205078125 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000000000126217744835361888865732358691622310176166669409751499176025390625 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000000000000631088724176809444328661793458111550880833347048757495880126953125 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000000000003155443620884047221643308967290557754404166735243787479400634765625 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000000000015777218104420236108216544836452788772020833676218937397003173828125 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000000000000078886090522101180541082724182263943860104168381094686985015869140625 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000000000000394430452610505902705413620911319719300520841905473434925079345703125 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000000000001972152263052529513527068104556598596502604209527367174625396728515625 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000000000000009860761315262647567635340522777992982513021047636835873126983642578125 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000000000000049303806576313237838176702613889964912565105238184179365634918212890625 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000000000000246519032881566189190883513069449824562825526190920896828174591064453125 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000000000000001232595164407830945954417565347249122814127630954604484140872955322265625 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000000000000006162975822039154729772087826736245614070638154773022420704364776611328125 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000000000000030814879110195773648860439133681228070353190773865112103521823883056640625 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000000000000000154074395550978868244302195668406140351765953869325560517609119415283203125 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000000000000000770371977754894341221510978334203070175829769346627780288045597076416015625 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000000000000003851859888774471706107554891671015350879148846733138901440227985382080078125 percent	-	45	42	38	-
Yeast extract 0.000000000000000000000000000000000019259299443872358530537774458355076754395744233665694507201139926910400390625 percent	-	45	42	38	-
Yeast extract 0.0000000000000000000000000000000000096296497219361792652688872291677537771978721168328472536005699634552001953125 percent	-	45	42	38	-
Yeast extract 0.00000000000000000000000000000000000481482486096808963263444361458387688859893605841642362680028498172760009765625 percent	-				

Table 1

C. botulinum 62A Spore Recoveries in Various Agar Media¹

Basal Medium	1 Per Cent Soluble Starch	Incubation, Days			
		3	6	17	33
Yesair's plus 0.1 percent starch	-	45	41	36 ²	-
Difco nutrient	-	28	30	22	-
Difco nutrient	+	35	42	35	-
Difco brain heart infusion	-	4	6	6	-
Difco brain heart infusion	+	12	23	30	30
BBL anaerobic	-	0	0	2	-
BBL anaerobic	+	0	1	6	8
Difco yeast extract, 1.0 percent	-	19	17	15	-
Difco yeast extract, 1.0 per cent	+	15	12	11	-
Corn steep liquor solids, 1.0 per cent	-	8	10	8	-
Corn steep liquor solids, 1.0 percent	+	7	9	11	14
Fresh beef infusion ³	-	43	50	47	-
Fresh beef infusion	+	60	60	53	-
Fresh liver infusion ³	-	24	24	23	-
Fresh liver infusion	+	46	52	49	-

¹ All media contained 0.2 per cent glucose and 0.2 per cent thioglycolate.

² The somewhat lower counts in this column as compared to the 6-day values is due to the obscuring of small colonies by development of gas bubbles and cloudiness in the media.

³ Prepared exactly like Yesair's medium (cf. Wynne and Foster, 1948.)

References

- Curran, H. R., and Evans, F. R. 1937 The importance of enrichments in the cultivation of bacterial spores previously exposed to lethal agencies. J. Bact., 34, 179-189.
- Foster, J. W., and Wynne, E. S. 1948 Physiological studies on spore germination, with special reference to Clostridium botulinum. IV. Inhibition of germination by unsaturated C₁₈ fatty acids. J. Bact., 55, In press.
- Morrison, E. W., and Rettger, L. F. 1930a Bacterial spores. I. A study in heat resistance and dormancy. J. Bact., 20, 299-311.
- Morrison, E. W., and Rettger, L. W. 1930b Bacterial spores. II. A study of bacterial spore germination in relation to environment. J. Bact., 20, 313-342.
- Olsen, A. M. and Scott, W. J. 1946. Influence of starch in media used for the detection of heated bacterial spores. Nature, 157, 337.
- Wynne, E. S., and Foster, J. W. 1948 Physiological studies on spore germination, with special reference to Clostridium botulinum. I. Development of a quantitative method. J. Bact., 55, In press.